

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/70, 35/78</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/06808</b> <b>(43) International Publication Date:</b> 27 February 1997 (27.02.97)
<b>(21) International Application Number:</b> PCT/FI96/00443 <b>(22) International Filing Date:</b> 16 August 1996 (16.08.96) <b>(30) Priority Data:</b> 516,769 18 August 1995 (18.08.95) US <b>(71) Applicant (for all designated States except US):</b> XYROFIN OY [FI/FI]; Kyllinkinportti 2, FIN-00240 Helsinki (FI). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> OLINGER, Philip, M. [US/US]; 1349 Linwood Drive, Terre Haute, IL 47802 (US). PEARSON, Julita [GB/GB]; 219 Pickhurst Rise, West Wickham, Kent BR4 0QA (GB). SAUNDERS, David [GB/GB]; Spring Hill, 12 Compton Way, Farnham, Surrey GU10 1QZ (GB). <b>(74) Agent:</b> OY KOLSTER AB; Iso Roobertinkatu 23, P.O. Box 148, FIN-00121 Helsinki (FI).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> COMBINATION OSMOTIC AND BULK FORMING LAXATIVES  <b>(57) Abstract</b>  A combination bulk-forming and osmotic laxative is provided by combining a bulk-forming laxative in the form of psyllium hydrophilic mucilloid with an osmotic laxative such as lactitol. The addition of the osmotic laxative to the bulk-forming laxative with a binder solution and subsequent drying produces an improved laxative in granulate form which is directly compressible into tablets.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## COMBINATION OSMOTIC AND BULK FORMING LAXATIVES

**Field of the invention:** The present invention relates generally to laxatives. Particularly, the present invention relates to a combination laxative which includes an osmotic laxative and a bulk-forming laxative. Still more particularly, the present invention relates to the granulated combination of an osmotic laxative in the form of a sugar alcohol, such as lactitol, with a bulk forming laxative such as psyllium to provide a laxative form which is directly compressible.

### BACKGROUND AND SUMMARY OF THE INVENTION

There are four general types of laxatives that are currently available: 1) bulk-forming; 2) osmotic; 3) stimulatory; 4) softening agents. The present application is directed toward the bulk-forming and osmotic types and specifically towards the use of an osmotic laxative in enhancing and correcting deficiencies found in bulk-forming laxatives.

While bulk-forming laxatives such as those formed from psyllium husks provide generally safe and effective laxatives, psyllium laxatives generally suffer from three primary deficiencies. First, bulk-forming laxatives such as psyllium are not directly compressible and lack the ability to be readily formulated in tableted form. Second, they do not disperse well and often form lumps of psyllium material, the interiors of which are still substantially dry. Third, psyllium seed husk has very poor wetting qualities and therefore does not mix well.

The present invention addresses the poor compressibility of psyllium husk by combining the ground psyllium husk with a sugar alcohol, such as lactitol, which, as an osmotic laxative, has its own laxative properties.

Specifically, psyllium hydrophilic mucilloid has not yet successfully been provided in a tablet form but

only in a powder form that may be dispersed in water. A chewable tablet containing psyllium would be beneficial because the masticating action in the mouth would aid dispersion. The present invention combines psyllium, which is not directly compressible, with a sugar alcohol to form a compressible granulate which may be compressed into tablets. Thus, the present invention provides a solution to problems inherent with one laxative -- bulk-forming laxatives such as psyllium seed husks -- by combining the bulk-forming laxative with another laxative -- an osmotic laxative, such as lactitol.

By way of background, bulk-forming laxatives rely upon the use of dietary fiber. Dietary fiber is the portion of plant materials that is resistant to digestion. There are generally two types of dietary fiber: insoluble and soluble. Insoluble dietary fiber passes through the digestive system in a substantially intact form. Insoluble fiber passes through the digestive system quickly because it does not absorb water, like soluble fiber.

In contrast, soluble fiber absorbs water in the small intestine and in the stomach. Psyllium is one source of soluble fiber. Specifically, the psyllium form known as psyllium hydrophilic mucilloid consists of the husk (either milled or unmilled) of blond psyllium seeds. When exposed to water in the stomach and small intestine, the psyllium hydrophilic mucilloid forms a gelatinous mass which aids in the treatment of constipation by acting as a fecal softening agent and also acts to soothe and protect the small intestine.

However, obtaining an appropriate gelatinous mass of psyllium hydrophilic mucilloid in the small intestine is problematic due to the poor dispersability and mixability of the psyllium hydrophilic mucilloid in water prior to ingestion. The individual particles of the psyllium hyd-

rophilic mucilloid tend to lump or agglomerate together when mixed with water. The psyllium hydrophilic mucilloid often fails to evenly disperse throughout the gelatinous mass and the hydration of psyllium hydrophilic mucilloid tends to take place over the exterior surface of such an agglomerated mass to form gelatin-coated lumps, the interiors of which are substantially dry. These gelatin-coated lumps are less voluminous than properly dispersed psyllium hydrophilic mucilloid and therefore the laxative benefits of the psyllium are compromised.

The poor dispersability of psyllium hydrophilic mucilloid is exacerbated by its poor mixing characteristics. Psyllium hydrophilic mucilloid is difficult to wet and tends to float on the surface of water. When mixing with water is attempted, the psyllium hydrophilic mucilloid forms lumps or masses. Psyllium is also difficult to mix when combined in the batter of a baked or continuously extruded food. Often, lumps of psyllium and material will result which result in an unacceptable food product due to textural deficiencies or to the poor taste of the psyllium material and psyllium derivatives.

In contrast to bulk-forming laxatives, osmotic laxatives act to retain water in the colonic lumen thereby counteracting the normal dehydrating action of the colon. By suppressing the dehydration action of the colon, the osmotic laxative produces a fecal stream which is softer, bulkier and easier to expel.

Lactitol may be used as an osmotic laxative because it is not absorbed in the small intestine and is thereafter hydrolysed and fermented in the large bowel by the action of intestinal microflora resulting in the liberation of galactose and sorbitol which are thereafter metabolized to organic acids and short chain fatty acids by colonic anaerobes. An osmotic effect is created in the co-

lon, which normally acts to hydrate the fecal stream. The reduction in the dehydrating action of the colon produces a softer, bulkier fecal stream. Studies have shown that administration of 10 to 20 grams of lactitol daily results in a predictable laxation response of typically 24 hours or less. In contrast, other sugar alcohols have not proven to be effective osmotic laxatives and are therefore not recognized as such because they are unpredictable and not necessarily dosedependent.

Additionally lactitol exhibits other properties which makes it a preferred candidate for combination with psyllium. Specifically, lactitol does not induce an increase in blood glucose or insulin levels and contributes half the calories of most other carbohydrates (2 calories per gram). Because the control of blood glucose, lipids and body weight are primary goals in diabetes management, lactitol is considered to be a beneficial product for diabetics. Lactitol is also non- cariogenic and has a moderate, sugar-like sweetness. Lactitol also has good solubility, low hygroscopicity, and when provided in a crystalline form, is suitable for dry mixing. Finally, lactitol is also available at a price that is competitive with other sugar alcohols.

Thus, the two types of laxatives discussed above, bulk-forming and osmotic, operate by two entirely different mechanisms. Bulk-forming laxatives absorb water and expand in the colonic lumen; in contrast, osmotic laxatives retain water in the colonic lumen by osmosis.

While the prior art teaches the use of both types of laxatives and further teaches the use of psyllium as a bulk- forming laxative and the use of lactitol as an osmotic laxative, the prior art does not teach a combination of psyllium and lactitol. Further, the prior art does not teach the use of an osmotic laxative, such as lactitol, to

enhance the compressibility, dispersability and mixability of bulk-forming laxatives, such as psyllium. One prior combination of psyllium and a sugar alcohol, specifically mannitol, is taught in U.S. Patent No. 5,320,847 to Valentine et al. However, Valentine teaches the use of mannitol only as a low-calorie sweetener or substitute for free sugar, not as an osmotic laxative. Mannitol is not proven or recognized as a therapeutic osmotic laxative. Valentine also does not attempt to provide a psyllium laxative in a tablet form.

A laxative provided in tablet form could be designed either for immediate swallowing or chewing. As noted above, psyllium is not available in tablet form because it is simply not compressible. However, when combined with a sugar alcohol, such as lactitol, psyllium can be provided in a tablet form due to the sufficient compressibility of the psyllium/lactitol combination. The masticating action of the mouth aids in the dispersion of the psyllium. Providing psyllium in a tablet form also provides a vehicle whereby other ingredients, such as medications, minerals and vitamins can be delivered with the laxative material. Finally, tablets are easily transported by the consumer and easily dispensed by the pharmacist.

Two criteria are used to evaluate the quality of a tablet. Those two criteria are crushing strength or hardness and friability. Hardness is a measure of a tablet's strength. Hardness is measured by determining the lateral breaking strength expressed in kilograms, Strong Cobb Units (S.C.U.) or Newtons (N) exerted on a single tablet at the moment of rupture. Typical hardness testers are manufactured by Key Instruments. In the context of chewable tablets, the tablet hardness must be greater than about 30N to be commercially useful.

On the other hand, friability is another measure of a tablet's ability to resist chipping and breaking during

shipment and handling. Friability is measured under standard conditions by weighing out a certain number of tablets (generally 20 or more) and placing them in a rotating plexiglass drum in which they are lifted during replicate revolutions by a radial louver and then dropped through the diameter of the drum. The tablets are then re-weighed and a percentage of the powder that has rubbed off is calculated. Friability of less than about 10% is considered to be acceptable for most chewable tablet contexts.

It is therefore an object of the present invention to provide an improved laxative by combining a bulk-forming laxative with an osmotic laxative.

Yet another object of the present invention is to provide an improved psyllium bulk-forming laxative by combining psyllium with an osmotic laxative.

Another object of the present invention is to provide an improved bulk-forming laxative/osmotic laxative combination which may be provided in tablet form or powder form.

It is still another object of the present invention to improve the compressibility, dispersability and mixability of psyllium husk powder by combining the psyllium husk powder with lactitol.

Yet another object of the present invention is to provide a means for producing a compressible psyllium/lactitol granulate using fluid bed granulation techniques.

Other objects and advantages of the invention will become apparent upon reading the summary, description of the preferred embodiments and the appended claims.

The above and other objects are accomplished by the present invention in which it has been discovered that a combination of lactitol as an osmotic laxative and psyllium as a bulk-forming laxative. The combination of the present invention results in an improved laxative with the



qualities of both an osmotic laxative and an enhanced bulk-forming laxative due to the enhanced compressibility, dispersability and mixability of the psyllium.

5 One preferred embodiment involves the combination of psyllium hydrophilic mucilloid and lactitol. Specifically, approximately equal parts of psyllium hydrophilic mucilloid and lactitol are dry blended. The resulting powder mixture may be mixed with water and taken orally. Further, a binder solution may be subsequently mixed with  
10 the psyllium and lactitol to produce a granulate. The granulate may thereafter be dried and compressed into tablets.

If the granulate is to be compressed into tablets, the preferred form of lactitol is lactitol monohydrate.  
15 Three known binder solutions are available, including 60% lactitol solution, 3% polyvinylpyrrolidone (PVP) solution and 10% gelatin (250 Bloom) solution. The addition of 10% by weight microcrystalline cellulose has been found to improve the hardness and friability of the tablets.

20 If the resulting lactitol psyllium mixture is to be provided in a powder form, three known forms of lactitol are available. Specifically, the lactitol may be provided in the form of anhydrous lactitol crystals, lactitol monohydrate or anhydrous lactitol. It is preferred that the  
25 lactitol monohydrate and anhydrous lactitol monohydrate be milled to a maximum particle size of 125 microns or less. The preferred form of psyllium is Roeper (60 mesh).

The present invention may also be provided in the  
30 form of a baked product, such as a cookie, cracker or cake.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is provided in two forms: powdered and granulate. The granulate may either be dis-  
35 persed in water prior to ingestion or may be directly

compressed into a tablet form. The preferred granulate form of the present invention includes a dry blend of approximately equal parts of psyllium powder and lactitol. A liquid binder solution is then added and the psyllium, lactitol and binder solution is mixed and dried. The preparation is then compressed into tablet form. The following examples are provided.

#### EXAMPLES 1-4

100 grams of lactitol monohydrate was dry-blended with 100 grams of psyllium powder. 64 grams of a 60% lactitol solution (w/w) was added to the dry mixture and the three components were mixed continuously until thoroughly wetted. 1% magnesium stearate was added as a tableting lubricant. The blend was then dried overnight at 60-C and passed through a 1000 micron sieve to yield a granulate. The granulate was blended with 1% magnesium stearate (w/w) and tablets were prepared on a Manesty 2C single punch press using a 15 mm diameter beveled edge flat faced punch. The compression setting was increased until an acceptable tablet was obtained. The tablet hardness or crushing strength was measured using a Key Instruments tablet hardness tester. An average of 10 samples were taken. The friability of the tablets was measured using a rotary friability tester. Ten tablets were dropped 100 times with the weight loss being recorded. The following results were obtained:

	<u>Exh. 1</u>
Average weight	1.051 g
30 Average thickness	4.7337 mm
Average hardness	31 N
Average friability	47.4%

The granulate of Example 1 was also combined with 10% by weight of microcrystalline cellulose (MCC) prior to

tableting. As seen below, the addition of MCC increased tablet hardness and decreased tablet friability.

		<u>Exh. 2</u>
5	Average weight	1.0624 g
	Average thickness	4.7651 mm
	Average hardness	38 N
	Average friability	7.6%

10                Examples 3 and 4 include the same composition as that shown in Example 1, i.e., without the addition of microcrystalline cellulose. The following results were obtained:

15		<u>Ex. 3</u>	<u>Ex. 4</u>
	Average weight	1.2049 g	1.1944 g
	Average thickness	5.4505 mm	5.4139 mm
	Average hardness	35 N	31 N

20                EXAMPLES 5-7

Softer tablets were obtained by using polyvinylpyrrolidone (PVP) binder solution. Examples 5, 6 and 7 are presented below which illustrate the use of PVP binder solution. Again, 100 g 5lactitol was dry blended with 100 g psyllium powder. 29 g 3% PVP solution was added to the dry mixture and mixed until completely wetted. The mix was dried overnight at 60°C and sieved to obtain a suitable granulate. The granulate was blended with 1% magnesium stearate (w/w) and tabletted.

30		<u>Ex. 5</u>	<u>Ex. 6</u>	<u>Ex. 7</u>
	Average weight	1.0724 g	1.0676 g	1.0464 g
	Average thickness	5.116 mm	4.9942 mm	4.8462 mm
	Average hardness	11 N	16 N	18 N

EXAMPLES 8 AND 9

Tablets can also be formed from the mixtures bound with a gelatin binder solution. Again, 100 g lactitol and 100 g psyllium powder were dry blended. 35 g 10% gelatin (250 Bloom) solution was slowly added to the mixture and mixed until thoroughly wetted. The mix was then spread out onto a metal tray and dried overnight at 60°C and sieved to produce a suitable granulate. The granulate was blended with 1% magnesium stearate (w/w) and tabletted. The weight, thickness and hardness properties are as follows:

	<u>Ex. 8</u>	<u>Ex. 9</u>
Average weight	1.0723 g	1.0702 g
Average thickness	4.8415 mm	4.8365 mm
Average hardness	22 N	20 N

EXAMPLES 10-12

Approximately 100 g lactitol monohydrate which was milled and screened through a 125 micron sieve was added to 100 g psyllium powder. No binder solution was added to the mixture. The blend was mixed for 2 minutes with 1% added magnesium stearate and pressed into a tablets. The following results were obtained:

	<u>Ex. 10</u>	<u>Ex. 11</u>	<u>Ex. 12</u>
Average weight	1.1137 g	1.1208 g	1.1416 g
Average thickness	5.3001 mm	5.3633 mm	5.4582 mm
Average hardness	17 N	17 N	16 N

EXAMPLES 13 AND 14

Using the same procedure, i.e. no binder solution, approximately 100 g milled anhydrous lactitol and 100 g psyllium powder were combined and blended with 1% magnesium stearate for 1-2 minutes before being pressed into tablet form. The following results were obtained:

	<u>Ex. 13</u>	<u>Ex. 14</u>
Average weight	1.1384 g	1.1389 g
Average thickness	5.3301 mm	5.3198 mm
Average hardness	20 N	21 N

5

EXAMPLE 15

Finally, 100 g anhydrous lactitol crystals and 100 g psyllium powder were combined and blended for 1-2 minutes with 1% magnesium stearate. However, this mixture could not be compressed into tablet form and therefore must be provided to the consumer in powdered form.

Accordingly, if the product is to be provided in tablet form, a lactitol solution was found to be the superior binding solution and the addition of microcrystalline cellulose (MCC) was found to improve hardness and friability characteristics. If the product is to be provided to the consumer in powder form, the addition of a binder solution is not necessary.

Either the granulate form (Examples 1-9) or the dry mixture (Examples 10-15) can be used to manufacture laxative-containing baked goods. Two cookie recipes are as follows:

EXHIBIT 16

	lactitol monohydrate	9.2%
25	lactitol/psyllium granulate	
	(Example 1)	35.8%
	flour	18.3%
	shortening	17.5%
	eggs	10.4%
30	skim milk powder	7.5%
	salt	0.2%
	baking soda	0.3%
	cream of tartar	0.2%
	Acesulfame K	0.05%
35	flavor	0.5%

EXHIBIT 17

	lactitol monohydrate	30.0%
	psyllium	15.0%
	flour	18.3%
5	shortening	17.5%
	eggs	10.4%
	skim milk powder	7.5%
	salt	0.2%
	baking soda	0.3%
10	cream of tartar	0.2%
	Acesulfame K	0.05%
	flavor	0.5%

15 Thus, the present invention may be provided in a powder form, a granulate form, a tablet form and in a baked product. Additional components such as medications, vitamins and minerals may also be provided with the combination osmotic/bulk-forming laxative of the present invention. Intense sweeteners such as Acesulfame K and aspartame may be added to enhance the flavor of the products. Flavorings and colors may also be used. Other tabletting lubricants in addition to magnesium stearate will be apparent to those skilled in the art.

20 Although only 17 embodiments of the present invention have been illustrated and described, it will at once be apparent to those skilled in the art that variations may be made within the spirit and scope of the invention. Accordingly, it is intended 11 I that the scope of the invention be limited solely by the scope of the hereafter  
30 appended claims and not by any specific wording in the foregoing description.

WHAT IS CLAIMED IS:

1. A combination osmotic and bulk-forming laxative comprising:

5           lactitol in an amount from about 40% to about 60%,  
and       psyllium hydrophilic mucilloid powder in an amount  
from about 40% to about 60%.

2. The laxative of claim 1, wherein the lactitol  
and psyllium hydrophilic mucilloid powder are combined by  
10       dry blending.

3. The laxative of claim 1, wherein the lactitol  
and psyllium hydrophilic mucilloid powder are combined in  
a granulate form.

4. The laxative of claim 1, wherein the lactitol is  
15       further characterized as lactitol monohydrate particles in  
a milled form and having a particle diameter of less than  
125 microns.

5. The laxative of claim 1, wherein the lactitol is  
further characterized as anhydrous lactitol crystals in a  
20       milled form and having a particle size of less than 125  
microns.

6. The laxative of claim 1, wherein the psyllium  
mucilloid powder is further characterized as having a par-  
ticle size of less than 125 microns.

7. The laxative of claim 4, wherein the combination  
is further characterized as being a granulate which is  
25       directly compressible into tablets.

8. The laxative of claim 5, wherein the combination  
is further characterized as being a granulate which is  
30       directly compressible into tablets.

9. A process for producing a combination bulk-for-  
ming and osmotic laxative in granulate form, the method  
comprising the following steps:

dry blending approximately equal parts of psyllium  
35       hydrophilic mucilloid powder and lactitol,

adding a binder solution,

mixing the binder solution, psyllium hydrophilic mucilloid powder and lactitol together to form a wet granulate, drying the wet granulate to provide a dry granulate, screening the dry granulate to a desired particle size or distribution.

10. The process of claim 9, wherein the lactitol is further characterized as lactitol monohydrate particles having a diameter of essentially less than about 125 microns.

11. The process of claim 9, wherein the lactitol is further characterized as anhydrous lactitol particles having a diameter of essentially less than 125 about microns.

12. The process of claim 9, wherein the binder solution is further characterized as being aqueous lactitol (60%) solution.

13. The process of claim 9, wherein the binder solution is further characterized as being aqueous PVP (3%) solution.

14. The process of claim 9, wherein the binder solution is further characterized as being aqueous gelatin (10%) solution.

15. The process of claim 9 further comprising the following step:

compressing portions of the dried granulate into tablets.

16. A combination osmotic and bulk-forming laxative provided in tablet form, the laxative comprising:

a directly compressible granulate including lactitol and psyllium hydrophilic mucilloid,  
an intense sweetener, microcrystalline cellulose,  
a tableting lubricant.

17. The laxative of claim 16, wherein the intense sweetener is aspartame.



18. The laxative of claim 16, wherein the intense sweetener is Acesulfame K.

5 19. The laxative of claim 16, wherein the directly compressible granulate is further characterized as including lactitol in an amount from about 40 to about 60% and psyllium hydrophilic mucilloid powder in an amount from about 60% to about 40%.

10 20. The laxative of claim 16, wherein the directly compressible granulate is present in an amount ranging from about 60% to about 90% and the microcrystalline cellulose is present in an amount ranging from about 40% to about 10%.

15 21. The laxative of claim 16, further comprising therapeutic amounts of vitamins.

22. The laxative of claim 16, further comprising therapeutic amounts of minerals.

23. A process for producing a combination bulk-forming and osmotic laxative in tablet form, the method comprising the following steps:

20 dry blending a directly compressible granulate with microcrystalline cellulose and a tableting lubricant,

the directly compressible granulate comprising psyllium hydrophilic mucilloid powder in an amount ranging from about 40% to about 60% and lactitol in an amount ranging from about 60% to about 40%,

25 compressing the granulate to form individual tablets.

30 24. The process of claim 23, wherein the microcrystalline cellulose is present an amount ranging from about 40% to about 10%,

the directly compressible granulate is present in an amount ranging from about 60% to about 90%.

35 25. The process of claim 24 further comprising the following step between the dry blending step and compressing step:

adding in intense sweetener.

26. The process of claim 25, wherein the intense  
sweetener is aspartame.

5 27. The process of claim 25, wherein the intense  
sweetener is Acesulfame K.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 96/00443

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A61K 31/70, A61K 35/78

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4999200 A (ANGEL CASILLAN), 12 March 1991 (12.03.91)  --	1-27
A	EP 0362926 A1 (THE PROCTER & GAMBLE COMPANY), 11 April 1990 (11.04.90)  --	1-27
A	EP 0101891 A1 (G.D. SEARLE & CO.), 7 March 1984 (07.03.84)  -- -----	1-27

☐ Further documents are listed in the continuation of Box C.
 ☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

18 November 1996

Date of mailing of the international search report

25 - 11 - 1996

 Name and mailing address of the ISA/  
 Swedish Patent Office  
 Box 5055, S-102 42 STOCKHOLM  
 Facsimile No. +46 8 666 02 86

Authorized officer

 Eva Johansson  
 Telephone No. +46 8 782 25 00

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

28/10/96

International application No.  
PCT/FI 96/00443

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4999200	12/03/91	NONE	
EP-A1- 0362926	11/04/90	SE-T3- 0362926	
		AU-B- 636304	29/04/93
		AU-A- 4245289	05/04/90
		CA-A- 1331535	23/08/94
		DE-T- 68906746	28/10/93
		ES-T- 2055014	16/08/94
		HK-A- 99296	14/06/96
		IE-B- 62926	08/03/95
		JP-A- 2200167	08/08/90
		PT-B- 91852	09/08/95
		US-A- 5149541	22/09/92
		US-A- 5445831	29/08/95
EP-A1- 0101891	07/03/84	SE-T3- 0101891	
		AU-B- 554250	14/08/86
		AU-A- 1713983	26/01/84
		CA-A- 1206417	24/06/86
		JP-B- 8022818	06/03/96
		JP-A- 59033215	23/02/84
		US-A- 4459280	10/07/84
		US-A- 4548806	22/10/85